

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BJ

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/565	A1	(11) International Publication Number: WO 95/12402 (43) International Publication Date: 11 May 1995 (11.05.95)
(21) International Application Number: PCT/US94/12782 (22) International Filing Date: 7 November 1994 (07.11.94) (30) Priority Data: 08/149,175 5 November 1993 (05.11.93) US 08/318,042 4 October 1994 (04.10.94) US (71) Applicant: UNIVERSITY OF FLORIDA RESEARCH FOUNDATION, INCORPORATED [US/US]; 223 Grinter Hall, Gainesville, FL 32611 (US). (72) Inventors: SIMPKINS, James, W.; 1722 NW 11th Road, Gainesville, FL 32604 (US). SINGH, Meharvan; 3316 SW 41st Place, Gainesville, FL 32608 (US). BISHOP, Jean; P.O. Box 11678, Jacksonville, FL 32239 (US). (74) Agents: SUNSTEIN, Bruce, D. et al.; Bromberg & Sunstein, 11th floor, 125 Summer Street, Boston, MA 02110-1618 (US).		(81) Designated States: AU, CA, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ESTROGEN COMPOSITIONS AND METHOD FOR NEUROPROTECTION		
(57) Abstract The use of an estrogen compound is described for protecting a population of nerve cells from death. The compound can be used in the treatment of a subject affected by a neurodegenerative disorder, in order to prevent the loss of neuronal activity.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

5

ESTROGEN COMPOSITIONS AND METHODS FOR NEUROPROTECTION

10

TECHNICAL FIELD

The present invention relates to methods for protecting cells in the central nervous system of subjects from cell death and for stimulating neuronal survival through enhanced growth factor production.

BACKGROUND OF THE INVENTION

Neurodegenerative diseases have a major impact on society. For example, approximately 3 to 4 million Americans are afflicted with a chronic neurodegenerative disease known as Alzheimer's disease. Other examples of chronic neurodegenerative diseases include diabetic peripheral neuropathy, multiple sclerosis, amyotrophic lateral sclerosis, Huntingdon's disease and Parkinson's disease. Not all neurodegenerative diseases are chronic. Some acute neurodegenerative diseases include stroke, schizophrenia, and epilepsy as well as hypoglycemia and trauma resulting in injury of the brain, peripheral nerves or spinal cord. There is a need for improved therapeutic agents and methods for reversing or retarding neuronal damage associated with each of these conditions.

Neurodegenerative diseases and aging are characterized by a wide range of symptoms which vary in severity and range from individual to individual. For example, Alzheimer's disease is characterized by symptoms such as depression,

- 2 -

aggression, impairment in short-term memory, impairment in intellectual ability, agitation, irritability and restlessness. Since estrogen becomes deficient in post-menopausal women, and since estrogen is believed to affect mood, some studies have been undertaken to assess the relief of behavioral symptoms associated with Alzheimer's disease. Unfortunately, those clinical trials that have been performed to establish the beneficial effect of estrogen on Alzheimer's disease have concluded that no statistically significant improvements in the disease course or symptoms resulted from the treatment. Fillet et al. 1986, Psychoneuroendocrinology 11:337-345; Honjo et al. 1989, Steroid Biochemistry 34:521-524. In one study where only 1 female and 1 male patient were studied and no statistics were available, a rapid reduction in symptoms of senile dementia was observed when estrogen was administered to the female patient in a cocktail of drugs together with chorionic gonadotrophin, a vasodilator and a non-steroidal anti-inflammatory agent after a period as short as one week (Aroonsakul 1990, US patent 4,897,389). There is a need for a better understanding of the underlying process of neurodegeneration such that improved treatment protocols and effective drugs may be designed that are effective at treating the disease itself so as to bring about a longterm meaningful reversal of symptoms.

A common feature of neurodegenerative disorders and the process of aging in animals is the progressive cell damage of neurons within the central nervous system (CNS) leading to loss of neuronal activity and cell death. This loss of activity has been correlated with adverse behavioral symptoms including memory loss and cognitive deficits. Therapeutic agents that have been developed to retard loss of neuronal activity either have toxic side effects or are prevented from reaching their target site because of their inability to cross the blood-brain barrier. The blood-brain barrier is a complex of morphological and enzymatic components that retards the passage of both large and

- 3 -

charged small molecules thereby limiting access to cells of the brain. There is a need for novel therapeutic agents that are readily transported across the blood-brain barrier as well as for novel methods of treatment of

5 neurodegenerative disorders that directly target the damaged site and are non-toxic.

Traditional methods of treating neurological symptoms focus on; modifying the electrical impulse itself as it moves between and along neurons; or modifying the release or

10 degradation of neurotransmitters. It is now recognized that neuronal cell density has an important impact on function. In various pathological conditions, loss of cell density has been observed resulting from accelerated neuronal cell death. The pattern of degeneration of neurons typically

15 originates from the nerve terminals and progresses "backward" toward the cell body (retrograde degeneration). In several systems, lesioning of certain brain regions results in compensatory sprouting of axons. This plasticity of neurons is attributed at least in part to the presence of

20 trophic growth factors.

These findings have spurred efforts to identify therapeutic agents that compensate for cell loss by stimulating sprouting of dendrites and axons of remaining cells so as to improve the structural integrity of the

25 damaged region. However, the optimal density of neurons and neuronal extensions is a delicate balance between deficiency and excess, a balance that varies with the environment of the cells. This balance can be disrupted when therapeutic agents act on normal or inappropriate tissue. There is a

30 need therefore to target therapeutic agents at a therapeutic dose specifically to those regions where they are required, or, alternatively, to identify agents that have a natural specificity for the target site only.

To date, there are no safe and effective methods for

35 treating loss of neuronal activity. However, considerable attention has recently been focused on naturally occurring proteins, collectively called neurotrophic factors (see

- 4 -

Table I), that promote growth and maintenance of cells of the central nervous system (CNS) and sympathetic and sensory neurons of the peripheral nervous system. In particular, the administration of nerve growth factor (NGF), a protein which is normally transported retrogradely in the intact brain from the hippocampus to the septal cholinergic cell bodies as well as from the cortex to the nucleus basalis, provides trophic support to cholinergic neurons and has been shown in animal models to have utility in reducing the effects of neurodegeneration due to trauma, disease or aging. The septum and the nucleus basalis are part of a region of the brain known as the basal forebrain. The effectiveness of administering NGF in response to damage is supported by experiments that demonstrate that cholinergic neurons in the medial septum can be protected from retrograde degeneration by chronic infusion of exogenous NGF (Rosenberg et al. 1988, Science 242:1575-1578). Indeed, infusion of NGF has been shown to significantly attenuate retrograde degeneration of cholinergic neurons after transection of their connections in the fimbria (the septo-hippocampal pathway).

One of the major problems confronting the use of NGF as a therapeutic agent is finding an appropriate method of increasing the levels of NGF at the appropriate target site. NGF is a large molecule and as such cannot normally pass across the blood-brain barrier and therefore has very limited access to the cells of the brain. Invasive methods are commonly used to place externally administered NGF within the brain. These methods are not sufficient to target NGF specifically to those cells where it is required. Non-localized targeting not only decreases the amount of protein available at the target site but also results in stimulation of growth of neurons at inappropriate sites resulting in potential harmful effects for the subject.

Another disadvantage of administering NGF as a therapeutic agent is that of induction of an immunological response to this protein. There is a need therefore for a

- 5 -

compound that does not in itself cause an immune response but could stimulate the production of endogenous NGF.

Current methods for administering nerve growth factor across the blood-brain barrier include: polymeric implants, osmotic minipumps, cell therapy using genetically engineered autologous or heterologous cells secreting NGF for implantation into the brain, and methods of increasing the permeability of the blood-brain barrier thereby allowing diffusion of these molecules to cells in the brain. Where exogenous NGF is used, a relatively large amount of relatively costly recombinant protein is required.

Rather than these aforestated solutions to delivery of proteins, it would be desirable to: avoid invasive techniques; to control the amount and the site of delivery of neurotrophic proteins to sites where they are most needed thereby minimizing toxic side effects; and to minimize the health care costs of treatment.

An additional approach to treating neurological symptoms has followed the observation that certain amino acids (glutamic acid and aspartic acid) act as excitatory neurotransmitters that bind the N-methyl D-aspartate (NMDA) receptor. Excess release of these amino acids (EAA) causes overstimulation of the neurons in neurodegenerative diseases as well as in conditions of hypoglycemia or trauma, resulting in neuronal loss and behavioral dysfunctions. NMDA is a potent and toxic analogue of glutamate which has been shown in animal studies to mediate much of the neuronal death associated with head trauma, hypoglycemia, anoxia, hypoxia and other conditions, and compromises the flow of blood, oxygen or glucose to the central nervous system.


A number of synthetic compounds that act as antagonists of the receptor have been described and tested in animal models. The possibility that these compounds are toxic in humans remains unresolved. Despite many years of clinical research, these antagonists are not as yet available as therapeutic products for treating patients.

- 6 -

For the foregoing reasons there is a need for methods of protecting neurons from accelerated cell death caused by trauma or disease or by the aging process or by combinations of these factors. There is also a need for methods that
5 stimulate the production of neurotrophic growth factors using small molecules that are capable of crossing the blood-brain barrier and that have minimal side effects.

SUMMARY

A preferred embodiment of the invention is directed
10 toward a method of protecting a population of nerve cells from death that includes administering to a nerve cell population in an animal subject an effective dosage of an estrogen compound sufficient to cause the nerve cell
15 population to be protected from progressive cell damage leading to the death of the cells otherwise occurring without any intervention, wherein the estrogen has a general structure:



25 a tautomer thereof, or a pharmaceutically acceptable salt thereof

A further embodiment of the invention is directed toward a method of protecting a population of nerve cells from death, that includes administering an estrogen compound
30 in an effective dosage sufficient to cause the nerve cell population to be protected from progressive cell damage leading to death of the cells otherwise occurring without any intervention.

A further embodiment of the invention is a method of
35 treating a subject, undergoing loss of neuronal activity at a site within a subject, comprising administering an estrogen compound to the subject in an effective amount so

- 7 -

as to allow the compound to decrease the loss of neuronal activity.

A further embodiment of the invention is a method of increasing the amount of a naturally occurring neurotrophic factor available to a neurodegenerating site in CNS tissue, comprising administering an estrogen compound to neurons in a subject at a physiologically relevant dose so as to achieve a plasma level of the compound sufficient to cause an increase in the amount of the neurotrophic factor available to the neurons.

A further embodiment of the invention is a method for preventing a neurodegenerative disorder induced by overstimulation of excitatory amino acid receptors comprising; administering to a subject suffering from such neurodegenerative disorder, an effective amount of an estrogen compound.

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features will become better understood with regard to the following description, appended claims and accompanying drawings where:

Figure 1 shows a histogram of the effects of 17β -estradiol (E_2) on the age related release of lactate dehydrogenase (LD) in primary cortical neuronal cultures.

Figure 2 shows a histogram of the protective effects of E_2 on the cytotoxicity induced by various levels of hypoglycemia in C6 cells.

Figure 3 shows a histogram of the protective effects of E_2 on the cytotoxic effects of N-methyl D-aspartate (NMDA) in neuroblastoma cells.

Figure 4 shows active avoidance performance following 5 weeks of ovariectomy in intact, ovariectomized and E_2 replaced animals.

Figure 5 shows active avoidance performance following 28 weeks of ovariectomy in intact, ovariectomized and E_2 replaced animals.

- 8 -

Figure 6 shows the effect of 5 week ovariectomy and E₂ replacement on high affinity choline uptake in the frontal cortex of behaviorally naive rats.

Figure 7 shows the effect of 5 week ovariectomy and E₂ replacement on high affinity choline uptake in the hippocampus of behaviorally naive rats.

Figure 8 shows the protective effect of α -E₂ on SK-N-SH cell viability at 24 hrs. and 8(b) at 48 hrs.

Figure 9 shows examples of molecular structures for α and β estradiols capable of acting as neuroprotectants.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed toward methods for protecting a population of nerve cells in a subject from death and toward stimulating neurotrophic factors for protecting cells from cell death in an animal subject.

An "estrogen compound" is defined here and in the claims as any of the structures described in the 11th edition of "Steroids" from Steraloids Inc., Wilton N.H., here incorporated by reference. Included in this definition are non-steroidal estrogens described in the aforementioned reference. Other estrogen compounds included in this definition are estrogen derivatives, estrogen metabolites and estrogen precursors as well as those molecules capable of binding cell associated estrogen receptor as well as other molecules where the result of binding specifically triggers a characterized estrogen effect. Also included are mixtures of more than 1 estrogen, where examples of such mixtures are provided in Table II below. Examples of α estrogen structures having utility either alone or in combination with other agents are provided in Figure 9.

β estrogen is the β isomer of estrogen compounds. α estrogen is the α isomer of estrogen components. The term "estradiol" is either α or β estradiol unless specifically identified.

- 9 -

The term "E₂" is synonymous with β -estradiol 17 β -estradiol and β -E₂. α E₂, α -E₂, and α -estradiol is the α isomer of β -E₂ estradiol.

5 An "animal subject" is defined here and in the claims as a higher organism including humans having neurons subjected to forces that result in progressive cell damage and cell death.

"Neurotrophic growth factors" are defined here and in the claims as endogenous soluble proteins regulating
10 survival, growth, morphological plasticity or synthesis of proteins for differentiated function of neurons.

"Neurodegenerative disorder" is defined here and in the claims as a disorder in which progressive loss of neurons occurs either in the peripheral nervous system or in the
15 central nervous system. Examples of neurodegenerative disorders include: chronic neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Huntington's chorea, diabetic peripheral neuropathy, multiple sclerosis, amyotrophic lateral sclerosis; aging; and acute
20 neurodegenerative disorders including: stroke, traumatic brain injury, schizophrenia, peripheral nerve damage, hypoglycemia, spinal cord injury, epilepsy, and anoxia and hypoxia.

These examples are not meant to be comprehensive but
25 serve merely as an illustration of the term "neurodegenerative disorder."

- 10 -

TABLE I

**List of Characterized Proteins
Exhibiting Neurotrophic Activities**

5		
	Growth Factor	References*
	<hr/>	
	Proteins initially characterized as neurotrophic factors:	
10	Nerve growth factor (NGF)	Thoenen et al., 1987 Whittemore et al., 1987 Hafti et al., 1989
15	Brain-derived neurotrophic factor (BDNF)	Barde et al., 1982 Lei?? et al., 1989
	Neurotrophin-3 (NT-3)	Ernfors et al., 1990 Hohn et al., 1990 Maisonpierre et al., 1990 Rosenthal et al., 1990
20	Neurotrophin-4 (NT-4)	Hallbrook et al., 1991
	Neurotrophin-5 (NT-5)	Berkemeier et al., 1991
	Ciliary neurotrophic factor (CNTF)	Lin et al., 1989
25	Heparin-binding neurotrophic factor (HBNF)	Stockli et al., 1989 Kovesdi et al., 1990
	Growth factors with neurotrophic activity:	
30	Basic fibroblast growth factor (bFGF)	Morrison et al., 1986 Walicke, 1988
	Acidic fibroblast growth factor (aFGF)	Walicke, 1988
	Insulin-like growth factors (IGF's), insulin	Aizenman et al., 1986 Baskin et al., 1987
35	Epidermal growth factor (EGF)	Fallon et al., 1984 Morrison et al., 1987
	Transforming growth factor α (TGF α)	Derynck, 1988 Fallon et al., 1990
40	Interleukin 1	Spranger et al., 1990
	Interleukin 3	Kamegai, 1990
	Interleukin 6	Hama et al., 1989
	Protease ?? I and II	Monard, 1987 Olmsdorf et al., 1989 Whitson et al., 1989
45	Cholinergic neuronal differentiation factor	Yamamori et al., 1989

* References given refer to recent reviews or recent key publications.

50

- 11 -

TABLE II

Commercial Estrogen Preparations

- 5 1. Estrone Aqueous Suspensions Usual dosage IM, 0.1 to 0.5 mg, 2 to 3 times weekly.
2. Estradiol Estrase: 1 to 2 mg PO, daily for three weeks; one week off.
- 10 3. Estradiol Conjugates in Oil.
 - a. Estradiol-cypionate 1 to 5 mg IM; every 3 to 4 weeks-more than 20 preparations.
 - b. Estradiol-valerate 10 to 20 mg IM; every 4 weeks - more than 30 preparations.
- 15 4. Oral estrogen preparations 50 to 65 % estrone sulfate and 20 to 35 % equilin sulfate.
 - a. Premarin 0.3 to 2.5 mg PO daily for 3 weeks; one week off.
 - b. Estrocon 0.625 to 2.5 mg PO daily for 3 weeks; one week off.
 - c. Progens 0.625 to 2.5 mg PO daily for 3 weeks; one week off.
 - d. Many others 0.625 to 2.5 mg PO daily for 3 weeks; one week off.
- 20 5. Oral Esterified Estrogens 75 to 85 % estrone sulfate and 6 to 15 % equilin sulfate.
 - a. Estratab 0.3 to 2.5 mg PO daily for three weeks; one week off.
 - b. Menest 0.3 to 2.5 mg PO daily for three weeks; one week off.
- 25 6. Estropipate Piperazine Estrone Sulfate.
 - a. Ogen 0.625 to 5 mg PO daily for 3 weeks; one week off.
7. Ethinyl Estradiol.
 - a. Estinyl 0.02 to 0.5 mg PO daily for three weeks; one week off.
 - b. Feminone 0.05 mg PO daily for three weeks; one week off.
- 30 8. Quinestrol a fat stored, slow release form of ethinyl estradiol.
 - a. Estrovis 100 µg PO daily for 7 days; 100,ug weekly thereafter.
- 35 9. Diethylstilbestrol (0.2 to 0.5 mg PO daily for three weeks; one week off.
10. Chlorotrianisene Tace; 12 to 25 mg PO daily for three weeks; on week off.
- 40 11. Oral Estrogen-Antianxiety Agent Combinations Oral.
 - a. Milprem-200 or -400 (Conjugated estrogens and meprobamate).
 - b. PMB 20 or 400 (Conjugated estrogens and meprobamate).
 - c. Menrium 5-2, 5-4, or 10-4 esterified estrogens and chlordiazepoxide.
- 45 12. Estrogen and Androgen Combinations IM in Oil.
 - a. 13 preparations with 2 mg estradiol cypionate and 50 mg testosterone cypionate.
 - b. 12 preparations with 4 mg estradiol valerate and 90 mg testosterone enanthate.
 - c. 4 preparations with 8 mg estradiol valerate and 180 mg testosterone enanthate.
 - d. 6 preparations with various combinations of other estrogens and androgens.

50 Properties of estrogen.

Estrogen occurs in at least two isomeric forms, including β estrogen and α estrogen. β estrogens are pleiotrophic molecule with many biological activities. Clinical uses include treatment of osteoporosis, symptoms of

55 menopause and fertility control. In embodiments of the invention, β estrogen has also been shown to protect a subject against neuronal loss.

In comparison to β estrogen, α estrogen is typically believed to be at least 100-1000 times less potent in

- 12 -

estrogenic activity. Numerous examples have been reported in the literature that show that the biological effects of β estrogen are not shared by the α isomer. In fact, in the art, α estrogen is typically used as a negative control for β estradiol.

We have shown for the first time that α estrogen has a comparable activity to that of β estrogen for neuroprotection. The novel activity identified for α estrogen presents a number of advantages in the treatment of neurodegenerative diseases, trauma and aging. These advantages arise in situations which require treatment of males where the development of female traits is to be avoided and the treatment of females where the subject has increased susceptibility to endometrial, breast and cervical cancer.

In an embodiment of the invention, a novel use for estrogen has been identified, namely to cause arrest and/or reversal of progressive degeneration of neurons. The degenerative process ultimately leads to behavioral defects that accompany reduction in observed neuronal density. According to this invention, improvements in cognition, memory and other behavioral symptoms occur in subjects suffering from neurodegenerative disorders caused individually by disease, trauma or aging or a combination of these factors following administration of estrogen. Also according to this invention is an arrest in progression of a neurodegenerative disease such as by intervention and/or prevention of neuronal loss the patient will not develop or manifest a further decline in disease course. Evidence for this effect is shown in the accompanying examples and is correlated with biochemical effects identified both in vivo and in vitro.

In Example 1, in vitro studies have demonstrated cytoprotective properties of estrogen where comparatively low concentrations of α -E₂ and β -E₂ enhance viability in neuronal cells. In a further embodiment of the invention, pretreatment of cells with β -E₂ has been shown to protect

- 13 -

cells from death caused by glucose reduction or deprivation, a condition called hypoglycemia. Post-treatment with E₂ can rescue cells from the effects of hypoglycemia in a manner that dissipates with increasing length of hypoglycemia

5 (Example 2).

These examples demonstrate that both α and β estradiol at physiologically relevant doses exert a cytoprotective effect on both glial and neuroblastoma cells lines in vitro and that this cytoprotective effect can be distinguished
10 from a mitogenic action. While not wishing to be bound by theory, we hypothesize that estrogen exerts a direct protective effect on neuronal cells.

In vivo studies described in Examples 3-6 have shown that estrogen can reverse an impairment in non-spatial
15 learning. This impairment is correlated with a time dependent decline in choline acetyl transferase (ChAT) in both the frontal cortex and the hippocampus which is attenuated in animals treated with estradiol. The ChAT-containing nerve terminals in these two brain regions have
20 cell bodies located in the basal forebrain. In Example 3, rats have shown improvements in behavioral performances as determined by the active avoidance test following the addition of estrogen to estrogen deficient ovariectomized animals. Collectively, these data provide a method of
25 treating subjects through the modulation of basal forebrain cholinergic function by means of treating with estrogen so as to reduce loss of learning and memory associated with neuronal damage.

Examples 1-6 demonstrate a number of different pathways
30 through which estrogen compounds exert a cytoprotective effect on cells in the CNS. These include protection against hypoglycemia (Examples 2a, 2b, and 6) and protection from overstimulation of EAA (Examples 2c and 5), as well as by the stimulation of neurotrophic growth factor production
35 (Examples 4a and 4b).

Example 4a describes experiments demonstrating the increase in brain derived neurotrophic factor (BDNF)

- 14 -

determined by increased levels of BDNF mRNA while Example 4b describes the stimulation of nerve growth factor production determined by increased levels of NGF mRNA. A further cytoprotective effect of estrogen has been demonstrated by
5 the amelioration of the toxic effects of overstimulation of excitatory amino acid (EAA) receptors (Examples 2c and 5). Overstimulation of the EAA receptors has been identified as characteristic of a number of neurodegenerative disorders including epilepsy. In a further embodiment of the
10 invention, the estrogen compounds ameliorate the toxic effects of hypoglycemia that is a further cause of progressive cell damage leading to cell death (Examples 2a, 2b, and 6).

While not wishing to be bound by theory, it is
15 suggested that estrogen compounds act on a fundamental process that impacts cell viability and cell response to adverse conditions that result in damage and death and that this process underlies the observed phenomena. An example of such a mechanism includes the regulation of glucose to
20 cells.

These observations contrast with observations of the prior art. Although estrogen has been identified as having utility in treating adverse behavioral symptoms that accompany fluctuations in hormones associated with menopause
25 in aging women, the biochemical basis for these effects has never been determined. As such, the treatment of behavioral effects with estrogen in human subjects has been restricted to the treatment of menopause in women that demonstrate signs of deficiency in estrogen and use in prevention of the
30 sequelae of menopause, namely osteoporosis, corrected by replacement therapy of estrogen.

The human clinical studies do however demonstrate that externally administered estrogen is non toxic. In addition, these studies demonstrate that estrogen administered
35 intramuscularly subsequently reaches the brain as inferred by the behavioral effects of the treatment and as predicted from the structure of the molecule.

- 15 -

Although clinical studies by Sherwin 1988, Psychoneuroendocrinology 13:345-357, and Sherwin and Phillips 1990, Annals of the New York Academy of Sciences 592:474-5, have shown a general mood enhancing effect in
5 oophorectomized women following intramuscularly administration of estrogen at doses of 10 mg, the mechanism by which this effect occurred is unclear.

Clinical studies on women with Alzheimer's disease have shown acute benefit in some patients by improving mood and
10 reducing depression in response to short-term duration of estrogen treatment (Fillit et al. 1986, Neuroscience Abst. 12:945; Fillit et al. 1986, Psychoneuroendocrinology 11: 337-345; Honjo et al, 1989, Steroid Biochem. 34:521-525). The results from these studies were difficult to interpret
15 because of the small number of patients used, because treatment only had limited effect on a few patients, and because treatment extended for a short period of time only. Indeed neither study was of a duration long enough to evaluate or indeed to manifest any neuroprotective effects
20 if any of estrogen. An additional study by Aroonsakul, 1990, US Patent 4,897,389 described the effect of anabolic hormones including estrogen on a single man and a single
/ women where the relief of symptoms was proposed to result from stimulation of cell replication and human growth
25 hormone production following a short-term administration of hormone mixtures.

Biochemical studies on the action of estrogen on cells of the CNS either in vivo or in vitro has resulted in conflicting reports. A number of studies have shown that
30 estradiol has an effect on the plasticity of neurons. Morse et al. 1986, Experimental Neurology 94:649-658, reported that an estrogen derivative enhances sprouting of commissural-associational afferent fibers in the hippocampal dentate gyrus following entorhinal cortex lesions.
35 Additionally, cyclic changes in synaptic density in the CA1 of the hippocampus were shown to be related to circulating E₂ levels (Woolley et al. 1992, Journal of Neuroscience

- 16 -

12:2549-2554) and these changes could be mimicked with exogenous E₂ administration (Woolley et al. 1992). Indeed, it has further been shown that ovariectomy reduces and E₂ replacement normalizes high affinity choline uptake (HACU) in the frontal cortex of rats.

Additionally, Gibbs et al. 1993 (Society for Neuroscience Abstracts 19:5) have reported upregulation of choline acetyltransferase (ChAT) levels following estradiol treatment in the medial septum after 2 days and 2 weeks of treatment although no effect was observed after 1 week using in situ hybridization of ChAT mRNA. Luine et al. 1980, Brain Research 191:273-277, reported increased ChAT levels in the preoptic and hypothalamic regions of the rat brain in response to estradiol treatment.

In an embodiment of the invention, administration of estrogen in a physiological dose results in the reversal of impairment of non-spatial learning in female rats that have been ovariectomized (OVX). These behavioral effects of short-term OVX and E₂-replacement are correlated with biochemical changes in the hippocampus and the frontal cortex of the brain; in particular, a reduction and increase in high affinity choline uptake (HACU) in OVX and E₂-pellet treated rats, respectively. Short-term E₂-replacement also had a positive effect on choline acetyltransferase activity (ChAT) in the hippocampus, but not in the frontal cortex. Long-term E₂ replacement appeared to prevent the time-dependent decline of ChAT in the frontal cortex and to attenuate ChAT activity decline in the hippocampus. Collectively, these data show that estrogen has a cytoprotective effect on cells in the CNS and that the estrogen environment of adult female rats influences learning and the activity of basal forebrain cholinergic neurons and also demonstrate the importance of estrogens in the maintenance and proper function of basal forebrain cholinergic neurons in the female rat.

In 1977, Perez-Polo et al. (Life Sci. 21:1535-1543) published a paper entitled "Steroid Induction of Nerve

- 17 -

Growth Factor Synthesis in Cell Culture". Although the title indicates a positive effect of steroids on the production of NGF by glial cells in vitro, closer inspection of the paper revealed the opposite. By using non-specific polyclonal antibodies, increased levels of a high molecular weight molecule was identified in response to high levels of estrogen whereas there was no increase in the amount of a low molecular weight (MW) fraction now known to contain β NGF and similarly recognized by the antibody preparation.

5 The doses of estrogen used in the experiment were 5,500 times higher than physiological levels. The levels of the low MW fraction attributed to NGF and secreted into the media were found to be 1000-fold higher than predicted for NGF production using current methods. No further work was

10 carried out by these investigators or others to determine in vivo effects of estrogen on NGF production. Subsequent in vivo studies by Gibbs et al. 1993 reported that steroids (17- β estradiol) caused decreased levels of NGF mRNA in the hippocampus followed by decreased levels in the medial

15 septum and the diagonal band of Broca.

Contrary to the above report, an embodiment of the invention describes how estradiol stimulates the production of neurotrophic growth factor mRNA. For the first time, an estrogen compound has been described that not only has ready

25 access to brain tissue across the blood-brain barrier but can stimulate the availability of growth factor where it is most needed so as to reverse and offset the effects of neurodegeneration.

In a preferred embodiment of the invention, novel

30 properties have been identified for estrogen compounds that provide for the first time a method of using estrogen to arrest and/or reverse progression of neurodegenerative disorders rather than merely treating the symptoms of the disease.

35 In a preferred embodiment, both stereoisomers of estradiol, 17- β -estradiol and 17- α -estradiol have been found effective in reversing neurodegeneration.

- 18 -

In a preferred embodiment, estradiol is administered to rats and also to humans at concentrations sufficient to exert neuroprotective effects in the CNS. These doses vary according to interperson variability, the route of administration and the estrogen formulation used. For example, in rats, estradiol is administered subcutaneously by means of a silastic tube to achieve plasma levels of about 50 pg/ml. In humans, 0.2-10 mg or more specifically 1-2 mg of orally administered estrase (estradiol) given daily is commonly administered to patients suffering from post menopausal syndrome. These levels are also expected to be effective in treating neurodegenerative disorders in human subjects.

The recommended route of administration of the estrogen compound includes oral, intramuscular, transdermal, buccal, intravenous and subcutaneous. Methods of administering estrogen may be by dose or by controlled release vehicles.

Administration of estrogen may include the use of a single estrogen compound or a mixture of estrogens.

The protection of cholinergic neurons from severe degeneration is an important aspect of treatment for patients with acute or chronic neurodegenerative disorders, an example of chronic disease being Alzheimer's disease. For Alzheimer's patients, estrogen replacement or supplementation may be of significant therapeutic use. Other diseases for which estrogen treatment may be effective include Parkinson's disease, Huntington's disease, AIDS Dementia, Wernicke-Korsakoff's related dementia (alcohol induced dementia), age related dementia, age associated memory impairment, brain cell loss due to any of the following; head trauma, stroke, hypoglycemia, ischemia, anoxia, hypoxia, cerebral edema, arteriosclerosis, hematoma and epilepsy; spinal cord cell loss due to any of the conditions listed under brain cell loss; and peripheral neuropathy. Because of its cytoprotective properties, it is suggested that one pathway of action for estrogen is the inhibition of apoptosis.

- 19 -

Examples

5 **Example 1a:** In vitro studies demonstrate increased viability of nerve cells in the presence of an estrogen compound where estrogen exerts a cell protective effect.

Experimental design. A neuroblastoma (SK-N-SH) cell line was obtained from American Type Culture Collection (ATCC) (Rockville, MD). Culture conditions have been described previously (Keller et al. 1976, Kolbe et al. 1976). All experiments were accomplished in cells that were in passage number 3-6. Each experiment was divided into three treatment groups: Group 1 was RPMI media supplemented with 10% fetal bovine serum (FBS), group 2 was RPMI media without the FBS (serum free group) and group 3 was serum free RPMI media supplemented with 544 pg E₂/ml (E₂ provided by Steraloids, Inc., Wilton, NH), greatly in excess of the 16pg/ml of estrogen normally found in FBS. E₂ was initially dissolved in 100μl absolute ethanol and then diluted with media. Media for the other two groups were similarly prepared with the addition 100 μl absolute ethanol with the media. Media was changed at 48 h into the experiment to maintain the normal schedule of nutrient replenishment.

25 The rate of growth of the population of cells in each experimental group was measured.

Quantification of cell viability. Cell viability was assessed using the trypan blue dye exclusion method of Black et al. 1964, Exp. Cell Research 35:9-13. For each aliquot, two separate counts of total cells and dead cells were made. Live cell number was determined from the difference between total and dead cell number. An important limitation of the trypan blue method of staining is its time dependence. Therefore, care was taken to standardize time between resuspension of the cells, addition of the dye and actual counting on the hemacytometer.

Analysis of data. All data are presented as mean ± SEM after correcting for the dilution factor allowing the data

- 20 -

to be expressed as the number of cells per ml. Data were analyzed by an analysis of variance followed by Scheff 's F test. The criterion for significance was $p < 0.05$.

The mitosis ratio was calculated by dividing dead cells
5 (number cells/ml) over total cells (number cells/ml). This index provides a means of distinguishing between the mitotic and the cytoprotective effects of the treatments used in the experimental design. The ratio of dead/total cells was used to differentiate mitotic from cytoprotective effects of E_2
10 in culture.

Experimental results. The ratio of dead to total cells increased by 2 to 3 - fold with the removal of FBS from the culture media (Table III). The addition of E_2 prevented this serum removal effect and produced ratios similar to
15 those observed in the FBS group at each sample time. Through 48 h, the ratio of dead to live cells remained constant for E_2 -treated cells, but nearly doubled for serum free cells. By 96 h, the ratio increased in all three groups, indicating the diminishing cytoprotective effect
20 with time.

Although the total SK-N-SH cell number in the FBS group was approximately double that seen with E_2 treated cells at both 24 and 48 hours, the fraction of live cells was similar. However, E_2 -treated SK-N-SH cells did not show the
25 exponential growth pattern seen for cells grown in FBS. The ratio of dead to total SK-N-SH treated cells increased in all three groups with time (Table III). However, at each sampling time, this ratio was similar and markedly lower in both E_2 and FBS groups than under serum free conditions.

30

- 21 -

TABLE III
Dead/Total Cell Ratio for SK-N-SH Cultures

Time	Treatment	Sample Size (# Flasks)	Ratio (mean \pm sem)	Total Cells ($\times 10^4$ /ml)	Viable Cells ($\times 10^4$ /ml)	Dead Cells ($\times 10^4$ /ml)
10	24 hrs. Serum Free	6	$0.30 \pm 0.03^{\#}$	$19 \pm 2^{\#}$	$13 \pm 4^{\#}$	6 ± 1
	24 hrs. E ₂ (544 pg/ml)	6	0.16 ± 0.04	31 ± 5	26 ± 5	4 ± 1
	24 hrs. FBS	6	0.15 ± 0.03	34 ± 3	29 ± 3	5 ± 1
	48 hrs. Serum Free	6	$0.53 \pm 0.07^{\#}$	$21 \pm 3^{\#}$	$11 \pm 2^{\#}$	11 ± 2
15	48 hrs. E ₂ (544 pg/ml)	6	0.21 ± 0.05	41 ± 9	34 ± 9	7 ± 2
	48 hrs. FBS	6	0.14 ± 0.06	52 ± 7	46 ± 8	6 ± 2
	96 hrs. Serum Free	5	$0.62 \pm 0.12^{\#}$	$8 \pm 2^{\#}$	$4 \pm 2^{\#}$	4 ± 1
	96 hrs. E ₂ (544 pg/ml)	5	$0.38 \pm 0.06^*$	$22 \pm 4^*$	$14 \pm 2^*$	8 ± 2
20	96 hrs. FBS	5	0.22 ± 0.04	50 ± 8	40 ± 8	10 ± 1

[#] $p \leq 0.05$ vs both E₂ and FBS groups.

^{*} $p \leq 0.05$ vs both serum free and FBS groups.

25

The growth pattern of neuroblastoma cells in FBS-supplemented media revealed a doubling time of 6 to 9 days. A decrease in total number of cells with a corresponding increase in the fraction of dead cells was observed under serum free conditions across the five day evaluation.

Collectively, these data demonstrate that the enhanced total and live cell number observed results from a cytoprotective, rather than mitotic, effect of E₂ under specific in vitro conditions employed.

Example 1b. In vitro studies demonstrate increased viability of nerve cells in the presence of an estrogen compound where α -estrogen exerts a cell protective effect.

40

Experimental design. SK-N-SH cells were backcultured with 2 ml 0.02% EDTA (Sigma Chemical Corporation, St. Louis, MO), incubated for thirty minutes at 37°C and resuspended at a density of 1×10^5 cells per ml as follows. Each experiment had three treatment groups: Group 1 was RPMI media supplemented with 10% FBS (FBS group), group 2 was

- 22 -

RPMI media without the FBS (serum free group) and group 3 was serum free RPMI media supplemented with 544 pg/ml 17- α -E₂ pg/ml (α -E₂ group) (Steraloids, Inc., Wilton, NH). Treatment media were made the day of the experiment. α -E₂ was initially dissolved in 100 μ l absolute ethanol and then diluted with media. Media for the other two groups was similarly prepared with the addition and dilution of 100 ml absolute ethanol. The FBS has been assayed for β -E₂ determining concentrations of about 16 pg/ml. The RPMI 1640 media had undetectable β -E₂ levels. However, neither FBS nor media has been assayed for α -E₂.

Cell viability was assessed as described in Example 1a. After the cells had been incubated for 24 or 48 hours in the respective treatment media, cell suspensions were made by decanting media, topically washing each flask with 2 ml 0.02% EDTA, then incubating with 2 ml 0.02% EDTA for 30 minutes. Cells were subsequently resuspended in the appropriate media. 2 ml aliquots of five to six different flasks from each treatment were then treated with 500 μ l of 0.4% trypan blue stain (Sigma Chemical Corporation, St. Louis, MO). An important limitation of the trypan blue staining method for differentiating dead and live cells is its time dependence. Therefore, care was taken to standardize time between resuspension of the cells, addition of the dye, and actual counting on the hemacytometer.

Experimental results. As in Example 1a, the ratio of dead to total cells increased 2-3 fold with the removal of FBS from the culture media (Figure 8). The addition of α -estrogen prevented this serum removal effect and produced ratios similar to those observed in the FBS group at each sample time.

Example 1c. Estrogen delays or prevents time-dependent death of cells in primary cortical neuron cultures.

Experimental design. Primary cortical neurons were produced from rat brains that were 0-1 day old using a

- 23 -

variation of methods described by Monyer et al. 1989, Brain Research 483:347-354. Dispersed brain tissue was grown in DMEM/10% PDHS (pregnant donor horse serum) for three days and then treated with cytosine arabinoside (ARC) for two
5 days to remove contaminating glial cells. On day 5, the ARC media was removed and replaced with DMEM/10% PDHS. The neuronal cells were cultured for a further 4-7 days before use.

Control primary neuronal cultures show progressive
10 cell death between days 12 and 18 in culture. Twelve cultures were evaluated on days 12 and 16 for levels of the enzyme lactate dehydrogenase (LD) after adding estrogen to 6 cultures maintained in DMEM and 10% PDHS on day 9 and maintaining the remaining cultures as controls. LD was
15 assayed using a variation of the method by Wroblewski et al. 1955, Proc. Soc. Exp. Biol. Med. 90:210-213. LD is a cytosolic enzyme which is commonly used in both clinical and basic research to determine tissue viability. An increase in media LD is directly related to cell death.

20 **Results.** A single treatment on day 9 with estrogen significantly reduced ($p < 0.05$) the increase in LD observed in all 6 replicates on days 12 and 16. These data suggest that in primary neurons, estrogen exposure delays or prevents time-dependent death in culture for at least 7 days
25 (Figure 1), an observation that is further supported by examination of cultures by light microscopy. Here it was observed that estradiol prevented the retrograde degeneration (regression of neuronal extensions) and reduced the appearance of cytosolic inclusions (clusters of
30 material) in cell bodies; both of which are normally observed with aging in primary neuronal cultures in vitro and with degenerative disorders in vivo.

35 **Example 2a.** In vitro studies show that estrogen compounds protect cells against cyt toxicity induced by hypoglycemia.

- 24 -

Experimental approach. C6 glioma cells obtained from ATCC were plated in RPMI media with FBS at a concentration of 1×10^6 cells/ml in Falcon® 25 cm² tissue culture flasks. Four hours prior to the onset of hypoglycemia, the

5 maintenance media was discarded, monolayers were washed twice in the appropriate media and then incubated for four hours at 37°C in either serum free or serum free plus 544 pg/ml E₂. Kreb's Ringer Phosphate buffer was used to wash the monolayers twice before the addition of appropriate

10 glucose treatment. RPMI medium contains 2 mg glucose/ml; flasks were divided into groups of 6 each receiving 100% glucose (2 mg/ml), 80% glucose (1.6 mg/ml), 60% glucose (1.2 mg/ml) or 0% glucose (buffer) with no steroid addition or supplemented with 544 pg/ml E₂. All flasks were incubated

15 for 20 hours and then evaluated for total, live, and dead cell number utilizing the trypan blue method previously described.

Results. Figure 2 shows that hypoglycemia caused a marked and dose-dependent reduction in both total and live

20 C6 cell numbers in control flasks, which did not receive E₂ treatment. By contrast, at each of the levels of hypoglycemia tested, E₂ exposure prevented the loss in total and live cells associated with hypoglycemia. Trypan blue stains dying cells that have become permeable to the dye.

25 In adverse conditions, the number of cells in the total cell population that is measured is diminished as a result of disintegration of dead cells. Hence, the total cell numbers of cells in samples maintained for 20 hours in suboptimal levels of glucose show a reduction in total cell number in

30 Figure 2. However, a large percentage of this diminishing population are live cells. The addition of estradiol to cultures, maintained in suboptimal levels of glucose, protects the population from cell death and results in an overall greater number of live cells. The asterisk marks

35 those samples having a statistically significant reduction in total cells and live cells in the absence of estrogen when compared to samples in the presence of estrogen.

- 25 -

**Example 2b: Estr gen compounds protect cells
from cyt toxicity induced by
preexisting hypoglycemia.**

5 **Experimental design.** C6 cells were cultured as
described in Example 2a. On the day of the experiment,
flasks (5 to 6 per treatment group) were divided into 5
groups. One group was maintained in normal glucose media
(euglycemia: 2 mg/ml glucose in RPMI) and the remaining 4
10 groups were placed in RPMI with 80% of normal glucose
concentration (hypoglycemia). At one hour or at 4 hours
after the initiation of the hypoglycemia state, flasks were
treated with either RPMI (controls) or with E₂ (544 pg/ml
RPMI). At 24 hours after the initiation of hypoglycemia,
15 total, live and dead cells were counted as described above.
This study design permitted the determination of the
time-course of C6 cell rescued from the cytotoxic effects of
hypoglycemia.

Results. The results are shown in Table IV and
20 demonstrate the ability of E₂ to rescue C6 cells from the
cytotoxic effects of a preexisting hypoglycemia state. The
number of live cells was increased by more than 2-fold and
the number of dead cells was reduced by half after treatment
with E₂ at one hour after the initiation of hypoglycemia.
25 In contrast, treatment with E₂ at 4 hours after the
initiation of hypoglycemia had no effect on live cell number
and only slightly reduced dead cell number.

- 26 -

TABLE IV

Demonstration of the Ability of E₂ to Rescue C6 Cells from the Cytotoxic Effects of Hypoglycemia.

Glucose Pretreatment	N	Treatment	Time	C6 Cell Number (x10 ⁴ /ml)		
				Total	Live	Dead
Euglycemia	6		--	243±16	235±16	9±2
Hypoglycemia	6	SF - E ₂	1 hr.	108±7	19±3	89±6
	5	SF + E ₂	1 hr.	92±7	44±5*	49±4*
	6	SF - E ₂	4 hr.	124±10	17±2	107±8
	6	SF + E ₂	4 hr.	93±9	19±4	74±7

Depicted are mean ± SEM.

SF = Serum Free RPMI media.

* P < 0.05 from Serum Free group at 1 hour.

Example 2c: Estrogen compounds can protect cells against cytotoxicity induced by excitotoxic amino acids.

Experimental design. Five culture dishes containing SK-N-SH neuroblastoma cells were treated with E₂ (544 pg/ml) and 5 culture dishes were treated with RPMI media as described above. Four hours later, all cell were treated with NMDA (500 μM) for 5 minutes. Total live cells and dead cells were then determined.

Results. The protective effect of estradiol on the viability of the cell population is shown in Figure 3. Pretreatment with estradiol increased the number of live cells and reduced the number of dead cells in these neuroblastoma cultures following treatment with NMDA. These data demonstrate that E₂ pretreatment protects cells from the neurotoxicity associated with excitotoxic amino acids.

- 27 -

Example 3: In vivo studies show behavioral improvements caused by the neuroprotective effect of estrogen compounds.

5 The role of estradiol in protecting cholinergic function in the CNS and the association of this effect with learning and memory has been demonstrated as described below.

10 **Example 3a:** Learning and memory improve in estrogen treated ovariectomized rats.

Experimental design. Three groups of animals were analyzed using a standard 2-way active avoidance paradigm.
15 The three groups of animals were: ovary-intact, ovariectomized, and estradiol-replaced ovariectomized animals.

Animals. Young adult female (3-4 months old) CD-Sprague-Dawley rats (Charles River Breeding Laboratories,
20 Wilmington, MA) were maintained in standard breeding conditions.

Animal surgical procedure: Animals were anesthetized with methoxyflurane (Metofane, Pitman-Moore, Washington Crossing, NJ). Two-thirds of the rats underwent bilateral
25 ovariectomy using a dorsal approach. Three weeks following ovariectomy, a subset of the ovariectomized animals (the E₂ replaced group) received a 5 mm Silastic® (Dow Corning, Midland, MI) pellet containing a 1:1 mixture of cholesterol (Steraloids, Inc., Wilton, NH) and 17-B estradiol that was
30 implanted subcutaneously. Estradiol delivery through Silastic® tubing results from diffusion down a large concentration gradient and the fibrosis, which occurs over time around the Silastic® pellet, reduces diffusion. The E₂ replacement regimen was maintained for 2 or 25 weeks
35 following the 3 week rest period after ovariectomy. At 5 weeks and 28 weeks post ovariectomy, animals were behaviorally tested. In the long term treatment regimen, we removed and repositioned the Silastic pellets every 2 to 3

- 28 -

weeks to maintain E_2 diffusion from the Silastic. The ovariectomized group received sham pellets that were similarly re-positioned every 2 to 3 weeks. Both E_2 and sham pellets were washed twice with 100% ethanol and were
5 then incubated in PBS at room temperature for 48 hours prior to implantation. The resulting experimental groups were: ovary intact (INTACT), ovariectomized (OVX) (5 or 28 weeks), and estradiol-replaced (E_2 pellet).

Behavioral testing; active avoidance. To assess
10 learning, the 2-way active avoidance paradigm was employed following the procedures of Mouton et al. 1988, Brain Research 444:104-108. All three groups of animals were tested for 14 consecutive days, each day consisting of 15 trials. Each trial lasted for 1 minute; and consisted of the simultaneous presentation of conditional stimuli (a light and sound cue) for the first 5 seconds and a 7 second interval followed by an electrical foot-shock of 1.4 mA for a 2 second duration. Successful learning was determined by the number of correct responses or "avoidances" and was
20 defined as transferring from one side of the shuttle-box to the other within the first 12 seconds of each trial, before the onset of the footshock. In order to assess potential motivational differences between animals in different treatment groups, the number of "no transfers" were also
25 recorded. This parameter describes the number of trials in which the animal did not transfer from one side of the shuttle box to the other upon stimulation with the electrical shock.

Statistical analyses. Behavioral data were analyzed
30 non-parametrically using the Kruskal-Wallis one-way analysis of variance and the Mann Whitney U test for assessment of group differences.

Results. INTACT and E_2 -pellet animals performed better on the 2-way active avoidance paradigm relative to OVX
35 animals. Figures 4 and 5 show the average of the total number of avoidances made by each animal over the 14 day testing period starting at the end of the 5 week and 28 week

- 29 -

period in the three groups of animals. OVX rats showed a 59% decrease in the number of avoidances achieved relative to the intact group, but this difference was not significant. E₂ replacement of OVX rats caused a 8.5 - fold increase in the number of avoidances relative to the OVX group. When these same animals were maintained on their respective treatments and were behaviorally tested at 28 weeks, the number of total avoidances was increased in all groups relative to the 5 week testing period (Figures 4 and 5.) However, at this 28 week testing point, OVX significantly reduced the total number of avoidances by 61% and E₂ replacement continued to increase avoidances by 4.5-fold versus OVX rats. Furthermore, E₂-pellet rats showed a marked acceleration in their rate of learning at 28 weeks, achieving the criteria of performing correctly 11 out of a possible 15 times in a given day by 1.3 ± 0.3 days of testing (Table V). INTACT rats did not show this acceleration in learning requiring 9 ± 2.8 days to reach criteria. The OVX animals maintained their inability to learn the task in the allotted 14 days and were therefore assigned a value of 15 days.

At both 5 and 28 weeks, rats in each group maintained their relative order of proficiency in active avoidance and all animals performed better in the second trial than first (Figures 4 and 5). This enhanced performance during the second exposure to the paradigm likely reflects recall of the behavior learned during the first test. This long-term memory is particularly apparent in the E₂-pellet group, which at the 28 week time-point, reached the performance criteria in 1.3 ± 0.3 days (Table V). It appears that chronic exposure to low doses of E₂ may enhance long-term memory in addition to its stimulation of learning of this active avoidance paradigm.

- 30 -

TABLE V

Effect of Short-Term and Long-Term Ovariectomy and Estradiol Replacement on Learning and Retention

5	Treatment Group	Days to Reach Criteria	
		5 weeks (acute)	28 weeks (chronic)
10	INTACT	14.0 \pm 9.0	9.0 \pm 2.8
15	OVARECTOMIZED	15.0 \pm 0.0	15.0 \pm 0.0
	E ₂ PELLET	9.5 \pm 2.1 [#]	1.3 \pm 0.3*

[#] p \leq 0.05 vs Ovariectomized and Intact animals.

20 * p \leq 0.05 vs Ovariectomized animals using the Mann-Whitney U nonparametric statistic.

Example 3b: Neurochemical Assays demonstrate the neuroprotective effect of estrogen.

Biochemical tests were performed on enzymes normally produced by healthy cholinergic neurons to establish whether a cytoprotective effect could be detected in vivo and correlated with behavioral improvements.

Experimental design. Two assays were used to measure the viability of neurons. These assays were high affinity choline uptake (HACU) and choline acetyl transferase (ChAT) activity. HACU was conducted on tissues from both the frontal cortex and hippocampus of the rats. The effect of different lengths of time following ovariectomy or E₂ replacement on behavior (active avoidance behavior) and on the activity of cholinergic neurons (HACU and choline acetyl transferase (ChAT) activity) was measured.

Female Sprague-Dawley rats were either ovariectomized (OVX) only, or ovariectomized for 3 weeks followed by subcutaneous implantation of a silastic pellet containing 17- β -estradiol (E₂ pellet) resulting in a replacement of E₂ to physiological levels. Ovary intact animals served as a positive control. Active avoidance behavior and choline

- 31 -

acetyltransferase (ChAT) activity in the frontal cortex and hippocampus were assessed at 5 weeks and 28 weeks post ovariectomy while high affinity choline uptake (HACU) was measured only at the 5 week time point.

- 5 **Active Avoidance Test.** This test was carried out as described in Example 3a.

Biochemical Analyses.

- 10 **(a) Plasma estradiol assay.** Following behavioral testing, or in the case of the untested animals following the period of treatment, animals were decapitated and trunk blood was obtained. The blood was centrifuged at 13,500 x g for 1.5 minutes and resulting plasma was aliquoted into a separate tube for estradiol level determination at a later date. Plasma concentrations of E₂ were assayed by
- 15 radioimmunoassay (RIA) using commercial kits supplied by Diagnostic Product Corp. (Los Angeles, CA). The range of assay detectability was 20-3600 pg/ml. All samples were quantified in a single assay.

- 20 **(b) High Affinity Choline Uptake (HACU).** HACU was assessed in behaviorally naive animals. Following decapitation, brains were removed from the skull and placed on an ice-cooled surface. The hippocampus and frontal cortex were then dissected and immediately placed into ice cold 0.32 M sucrose buffer (0.32 M sucrose, 1.0 mM EDTA, 100
- 25 μM TRIS-HCl, pH = 7.4 at 4°C). Average wet weights for the tissue regions dissected were 0 mg and 110 mg for the frontal cortex and hippocampus, respectively. Tissue samples were then homogenized with a dounce homogenizer at 400 rpm. Homogenized samples were subsequently centrifuged
- 30 at 1000 x g for 8 minutes at 4°C. The supernatant (S1 fraction) was centrifuged at 30,000 x g for 15 minutes at 4°C. Following this spin, the supernatant was discarded and the resulting pellet (P2 fraction) was resuspended in 2 ml of cold, oxygenated Krebs buffer (139 mM NaCl, 5 mM KCl, 13
- 35 mM NaHCO₃, 1 mM MgCl₂, 1 mM NaH₂PO₄, 10 mM glucose, 1 mM CaCl₂; and was oxygenated for 15 minutes with 95% O₂/5% CO₂). High affinity choline uptake was determined in triplicate in

- 32 -

the presence of 1 μ M [3H]-choline (final specific activity: 4.5 Ci/mmol, New England Nuclear, Cambridge, MA). Non-specific uptake was estimated by adding 5 μ M hemicholinium-3 (Sigma Chemical Corp., St. Louis, MO); these 5 hemicholinium-3 values were subtracted from total counts to obtain high affinity values. Each reaction tube contained 200 μ l of P2 suspension. Unused tissue was stored at -30°C for subsequent determination of ChAT activity and protein levels. Analysis of protein in the P2 preparation was 10 conducted according to the method of Bradford using Coomassie Blue Dye (5).

HACU differences were analyzed using a t-test since INTACT versus OVX and OVX versus E₂-treated were evaluated only once each in two separate studies. Analysis of 15 variance (ANOVA) was employed for other neurochemical analyses. Multiple comparisons among groups were performed using Scheffe's post-hoc test.

(c) **ChAT assay.** ChAT activity was determined in both behaviorally naive animals (5 week group) as well as the 20 behaviorally tested group (28 week group). ChAT was assayed following a modified version of Fonnum 1975, J. Neurochem. 24:407-409. P2 samples were thawed and sonicated in the presence of 1% 1-butanol and centrifuged at 13000 x g for 5 min. 20 μ l of the resulting supernatant was used in each 25 reaction tube. The reaction mixture contained 0.28 mM [3H]-ACoA (specific activity: 45 μ Ci/mol, New England Nuclear, Cambridge, MA), 7.8 mM choline chloride and 0.2 mM physostigmine (Sigma Chemical Corp., St. Louis, MO). Incubation with the [3H]-ACoA was carried out for 30 min. 30 The reaction was terminated by the addition of ice cold glycyl-glycine buffer (GLY-GLY) at pH 8.6. Following a 10 minute incubation at 4°C, tetraphenyl boron dissolved in butyronitrile (10 mg/ml) was added to the reaction tube allowing for liquid cation exchange extraction of 35 acetylcholine (ACh). Samples were vortexed and centrifuged in a bucket centrifuge at low speed (185 x g) for 5 minutes to allow settling and separation of the organic and aqueous

- 33 -

phases. 100 μ l of the organic phase was then aliquoted into 7 ml scintillation vials and 4 ml of scintillation fluid (Liquiscint®, National Diagnostics, Atlanta, GA) were added. Vials were then counted in a Hewlett-Packard scintillation counter for 5 minutes each, the dpm were converted to pmoles and values were normalized for protein content.

Results.

(a) **Estradiol concentrations.** Serum E_2 concentrations were 43 ± 10 and 36 ± 5 pg/ml for the INTACT and E_2 -pellet groups, respectively. Ovariectomy reduced serum E_2 concentrations to below the sensitivity of the radioimmunoassay employed (20 pg/ml) in all but 5 animals sampled. These 5 animals, however, had serum levels that were very close to the sensitivity limit of the assay.

15 (b) **High Affinity Choline Uptake.** Ovariectomy significantly reduced HACU by 24% in the frontal cortex (Figure 6) and by 34% in the hippocampus (Figure 7). E_2 -replacement resulted in a reversal of this effect of ovariectomy, increasing HACU by 82% in the frontal cortex 20 (Figure 6) and by 46% in the hippocampus (Figure 7). Because HACU is a measure of cholinergic activity, it is concluded from these results that estradiol reverses the decline of cholinergic activity in the frontal cortex and hippocampus and further stimulates activity of cholinergic 25 projections to these regions.

(c) **Choline Acetyl Transferase activity.** In the frontal cortex, no significant differences in ChAT activity were detected within 5 weeks of ovariectomy (Table VI). At the 28 week time point, ChAT levels in the frontal cortex 30 were reduced in both the INTACT and OVX groups by 61% and 56%, respectively. In the E_2 -pellet group, however, this reduction was only 16%. In the hippocampus, five weeks of ovariectomy was a sufficient time period to induce a significant reduction in ChAT activity, and 3 weeks of 35 E_2 -replacement reversed this effect (Table VII). The reductions in hippocampal ChAT activities in INTACT and OVX animals between 5 and 28 weeks were comparable with those

- 34 -

seen in the frontal cortex. In E₂-pellet animals, loss of hippocampal ChAT activity was larger than that seen in the frontal cortex but was less than that seen in OVX or INTACT animals.

5

TABLE VI

Effect of Short Term and Long Term Ovariectomy
and Estradiol Replacement on Choline
Acetyltransferase Activity in the Frontal Cortex

10

Treatment Group	ChAT activity (nmol/30 min/mg protein)	
	5 weeks	28 weeks
INTACT	10.2 ± 0.5	4.0 ± 0.1
OVARECTOMIZED	9.2 ± 0.6	4.0 ± 0.2
E ₂ PELLET	9.8 ± 0.6	8.2 ± 0.8*

15

20

25

n = 6 for ovariectomized and E₂ pellet groups for the 5 week time period.
n = 5 for intact group for the 5 week time period.
n = 6 for all treatment groups for the 28 week time period.
* p ≤ 0.05 vs intact group and OVX group.

30

TABLE VII

Effect of Short Term and Long Term Ovariectomy and
Estradiol Replacement on Choline Acetyltransferase
Activity in the Hippocampus.

35

Treatment Group	ChAT activity (nmol/30 min/mg protein)	
	5 weeks	28 weeks
INTACT	13.2 ± 0.8	5.7 ± 0.3
OVARECTOMIZED	10.3 ± 0.3*	6.2 ± 1.1
E ₂ PELLET	12.7 ± 0.5	8.0 ± 1.1

40

45

50

n = 6 for ovariectomized and E₂ pellet groups for the 5 week time period.
n = 5 for intact group for the 5 week time period.
n = 6 for all treatment groups for the 28 week time period.
* p ≤ 0.05 vs. intact group and E₂ pellet group.

55

- 35 -

Example 4a: In situ hybridization demonstrates increased levels of brain derived neurotrophic factor (BDNF) mRNA is stimulated by estradiol.

5 This example demonstrates that estrogen can stimulate the production of neurotrophic growth factors such as BDNF.

Experimental design. Animals were ovariectomized for 12 weeks, while E₂ treated animals were ovariectomized for 3 weeks followed by 9 weeks of E₂ treatment. A second set of

10 animals were ovariectomized for 28 weeks and the E₂ treated animals were similarly ovariectomized for 3 weeks followed by 25 weeks of E₂ replacement. Intact control animals were run in parallel with each set of animals. The rats were deeply sedated with sodium pentobarbital. Immediately

15 following, the brains of these animals were transcardially perfused with 4% paraformaldehyde in 0.1M phosphate buffer. On the following day, the brains were removed from the skull and immersed in a series of solutions (cold 4% paraformaldehyde solution for 2 days at 4°C followed by a 4

20 % paraformaldehyde solution containing 20% (wt/vol) sucrose). After removal from the sucrose solution, the brains were blocked (removed olfactory bulbs, brainstem and cerebellum), frozen on dry ice and immediately stored at -80°C until time of use. 25 µm slices of the brain were

25 made using a microtome and placed in a 4% paraformaldehyde solution. Within 3 weeks, the slices were hybridized according to the method described by Gall et al. 1989, Science 245:758-761. The BDNF probe was 750 b.p. of rat BDNF cDNA, corresponding to the entire amino acid coding

30 region. The probes were labeled with ³⁵S using the T3 polymerase kit. The hybridized slices were mounted onto Vectabond treated slides and exposed to autoradiographic film for 4 days. The varying intensities of radioactive signal on each slice (the hybridization signal) was

35 translated into differing optical densities on film and was analyzed using the BRS 2 Imaging System (Imaging Research Inc.). The relative optical densities and the background levels were recorded. Signal was defined as the optical

- 36 -

density of the region divided by the average background level as assessed by parts of the film adjacent to the brain slice. In the imaging system employed, the higher the O.D., the lower the signal. As such, the data were transformed to the inverse of the signal to noise ratio in order to facilitate the presentation of the data.

Results. Ovariectomy resulted in a significant reduction in the BDNF signal in cortical sections relative to INTACT controls. Estradiol replacement of ovariectomized rats increased the BDNF signal to that normally observed in INTACT controls. The data in Table VIII derived multiple slices of the cerebral cortex of an animal in each treatment group demonstrate the stimulatory effect of an estrogen on BDNF synthesis.

TABLE VIII

Relative Level of BDNF mRNA in the Cortex of Intact, Ovariectomized and E₂ Pellet Animals

Treatment Group	BDNF Level
INTACT	1.242 ± 0.035
OVARECTOMIZED	1.159 ± 0.023*
E ₂ PELLETT	1.224 ± 0.026

* p ≤ 0.05 vs Intact and E₂ Pellet animals.

Example 4b: In vivo studies demonstrate increased levels of NGF mRNA stimulated by estradiol.

This example demonstrates that estrogen can stimulate the production of neurotrophic growth factors such as NGF. Similarly treated animals as in example 4a were used to measure levels of NGF mRNA using the dot blot technique. Animals were ovariectomized for 12 weeks, while E₂ treated animals were ovariectomized for 3 weeks followed by 9 weeks of E₂ treatment. RNA was isolated from the frontal cortex

- 37 -

and hippocampus using acid guanidinium isothiocyanate followed by phenol/chloroform extraction. The use of the dot blot technique was first validated by performing Northern blots to ensure that hybridization with our NGF probe of 771 b.p. recognizing the entire pre-pro sequence of NGF, resulting in a single band corresponding to the length of NGF mRNA. The NGF mRNA signal was normalized to the amount of RNA loaded as estimated by the amount of signal produced by subsequent actin hybridization.

Following 3 months of ovariectomy, NGF mRNA levels were significantly reduced (45%) in the frontal cortex (Table IX). E₂ treatment resulted in a partial recovery, albeit non-statistically significant from the OVX group. Hippocampal NGF mRNA was not found to differ from controls using this method of detection. E₂ treatment did, however, result in a significant increase in NGF mRNA levels (Table IX).

TABLE IX

The Effect of Three-Month Ovariectomy and Estrogen Replacement on NGF mRNA Levels in the Frontal Cortex and Hippocampus

Treatment	NGF mRNA (units per μ g actin)	
	Frontal Cortex	Hippocampus
INTACT	0.622 \pm 0.067 (n = 10)	0.590 \pm 0.052 (n = 6)
OVARECTOMIZED	0.342 \pm 0.050 [#] (n = 12)	0.616 \pm 0.051 (n = 12)
E ₂ REPLACED	0.453 \pm 0.070 (n = 8)	0.803 \pm 0.082* (n = 12)

[#] p \leq 0.05 vs Intact.
* p \leq 0.05 vs Ovariectomized.

- 38 -

Example 5: Proposed evaluation for the in vivo effects of estrogens in excitatory amino acid neurotoxicity.

5 Adult female rats are ovariectomized and two weeks later are treated with a "Silastic" pellet containing cholesterol (controls) or estradiol in amounts sufficient to elevate plasma estradiol levels in the physiologic range. After 1 to 2 weeks of such estrogen-replacement therapy, 10 rats receive an intracerebral injection of N-methyl-D-aspartate (NMDA) or artificial cerebrospinal fluid in amounts to induce extensive toxicity of brain neurons.

As an endpoint of the study, the number of neurons in the hippocampal CA1 region and the number of neurons in the 15 cerebral cortex are determined. In addition, nerve terminals in both regions would be stained for detection of choline acetyltransferase, the marker enzyme for cholinergic neurons.

It is expected that estrogen-replacement therapy will 20 reduce or eliminate the loss of nerve cells in both the hippocampus and in the cerebral cortex. Such a result teaches that physiological estrogen-replacement can protect brain cells from the neurotoxic effects of excitatory amino acids.

25

Example 6: Estrogen enhances cerebral glucose uptake at sites in the brain.

The effect of estradiol benzoate (E_2B) on cerebral 30 glucose uptake in various brain regions which contain variable numbers of E_2B receptors is described and the determination concerning increases in the amount of glucose transported into the brain in the presence of E_2B is also described for a selected animal model.

35 Female rats were bilaterally ovariectomized to eliminate endogenous estrogens and two to three weeks later were implanted with an atrial cannula for the i.v. administration of C^{14} -2-deoxyglucose (C^{14} -2-DG) to unanesthetized rats. Animals were allowed four to five days

- 39 -

to recover from the cannulation before the study was initiated.

On the day of the experiment, animals were randomized and assigned to groups which received either E₂B in oil or oil alone (controls) administered subcutaneously at the dose and times described below. In our initial study, animals were treated with oil or E₂B (10 µg/kg body weight) and were sacrificed at 2, 4, 8, 12 or 24 h. In an additional study, rats were treated with oil or E₂B at doses of 1, 10 or 100 µg/kg body weight and were sacrificed 4 h later. Forty-five minutes prior to sacrifice, all rats received a single injection via the atrial cannulae of C¹⁴-2-DG (25 µCi/ml saline/kg body weight; specific activity 49-53 mCi/mmol, New England Nuclear, Boston, MA).

To assess the effects of E₂B on transport of glucose across the blood-brain barrier we used the technique of Oldendorf (Oldendorf, Brain Res. 24:37-46; 1970; and Am. J. Physiol. 221:1629-1638 (1971)). One µCi/ml of C¹⁴-2-DG (specific activity 49-53mCi/mmol, New England Nuclear, Boston, MA) and approximately five µCi/ml of ³H₂O (specific activity 1 mCi/ml, New England Nuclear, Boston, MA) were mixed with Krebs' Ringer Phosphate solution buffered to pH 7.4 with 10mM HEPES and injected into the carotid artery of female rats.

Fifteen seconds after injection, the animals were killed by decapitation, trunk blood was collected for later assay of serum E₂B concentrations and the brain was removed from the cranium for dissection of the following regions: medial basal hypothalamus (MBH), preoptic area (POA), cortex, hippocampus, striatum, cerebellum and brainstem. The anterior pituitary (AP) was also isolated from the cranium. The dissection of brain tissues followed the methods of Glowinski and Iversen (Glowinski et al., J. Neurochem. 13:655-669 (1966)). Tissues were immediately weighed and placed in scintillation vials for processing. The mg weights (mean ± SEM) of tissues used in these studies were: MBH = 13.4±1.3; POA = 10.5±0.5; cortex = 38.7±3;

- 40 -

hippocampus = 27.7 ± 1.7 ; striatum = 24.3 ± 2 ; cerebellum = 32.5 ± 1.9 ; brainstem = 32.9 ± 1.9 and AP = 8.6 ± 0.6 .

During the procedure, peripheral plasma glucose values were in the normal range of 90-120 mg%.

5 Alternatively, half of the brain was dissected rostral to the midbrain and ipsilateral to the injection side, tissue passed through a 20 gauge needle and the sample subjected to routine digestion and then prepared for liquid scintillation counting as described above. An aliquot of
10 original isotope mixture was obtained by recovering the residual mixture in the injection syringes. Both aliquot and tissue samples were then counted for H^3 and C^{14} by routine liquid scintillation counting. Uptake by the Brain Uptake Index (BUI) was calculated after correcting for
15 counting efficiency using the following equation for extraction (E):

$$E = \frac{C^{14} \text{ in brain tissue} / ^3H \text{ in brain tissue}}{C^{14} \text{ in mixture} / ^3H \text{ in mixture}} \times 100$$

20 Table XI provides a sample of data generated within one experiment. Evaluations of the time or dose effect of E_2B on glucose uptake were done using a one way ANOVA. Post hoc comparisons were done with Dunnett's tests. These
25 statistical analyses were performed on raw data (dpms C^{14} /mg tissue) by comparing the dpms for the E_2B -treated group with its control (oil) group at each dose and at each time point evaluated. For clarity of presentation, the raw data were then expressed as percent of mean control. The magnitude of
30 the response of each E_2B -treated animal was determined by calculating the percentage increase of 2-DG uptake relative to the mean value of 2-DG uptake in the oil-treated control group. BUI data were analyzed by a one-tailed t-test for independent samples. Statistical difference was set at $p <$
35 0.05 for all tests.

The time course of the effects of a 10 μ g/kg body weight dose of E_2B on glucose uptake is shown in Table X. Overall, E_2B increased glucose uptake significantly in ovariectomized rats by 20 to 120% in 7 of 8 regions

- 41 -

examined. However, the time at which a significant increase was observed varied among regions. For 5 of 8 regions, the POA, hippocampus, striatum, cerebellum and AP, the peak E₂B effect was observed at 2 to 4 hours, while 2 regions showed peak effects at 12 hours (the MBH and cerebral cortex). Five regions showed a significant decline in glucose uptake at some time-point after an E₂B-induced increase. These regions were the POA, the hippocampus, the striatum, the cerebellum and the AP. No significant effect of E₂B on brainstem glucose uptake was observed.

The dose-dependency of the E₂B effect on brain glucose uptake at 4 hours post-E₂B injection is shown in Table XI. The 1 µg/kg dose of E₂B had no effect on brain glucose uptake. In contrast, the 10 µg/kg dose of E₂B increased glucose uptake in 6 of 8 regions examined at the 4 h time point. Furthermore, only two regions exhibited increased glucose uptake at the 100 µg/kg dose of E₂B, the MBH and the POA. All other regions examined exhibited a reduced glucose uptake, with three regions, the striatum, the cerebellum and the brainstem, showing a significant reduction.

Serum E₂ concentrations were observed to increase in a dose-dependent manner 4 h post E₂B injection. The 1 µg/kg E₂B dose did not elevate serum E₂B, levels above those observed in ovariectomized rats, the 10 µg/kg E₂B dose increased serum E₂B, levels into the physiological range, and the 100 µg/kg dose increased serum E₂B levels, which were 12 to 30 times those seen during peak serum E₂B concentrations on proestrus. Peak serum E₂B levels were observed at 4 hours after administration of a 10 µg/kg dose of E₂B.

Our evaluation of BUI using the Oldendorf method indicates that 4 h of exposure to E₂B increased the transport of glucose across the blood-brain barrier by about 40%. The 40% increase in BUI was accomplished by an E₂B-induced increase in C¹⁴-2-DG extraction with no change in ³H₂O extraction across the blood-brain barrier (Table XII).

- 42 -

It is expected that estrogen-replacement therapy will reduce or eliminate the loss of nerve cells in both the hippocampus and in the cerebral cortex. Such a result would indicate that physiological estrogen-replacement can protect
5 brain cells from the neurotoxic effects of hypoglycemia.

Although certain preferred embodiments of the present invention have been described, the spirit and scope of the invention is by no means restricted to what is described above. For example, while the examples describe experiments
10 performed in rats and cultured cells, it is believed that these are accurate models for predicting activity of compounds in humans.

- 43 -

TABLE X

Estrogen and Glucose Uptake in Various Brain Regions

5

Time Course Effects of a 10 μ g Dose of e_2 on
4 Hour Glucose Uptake in a Variety of Brain
Regions and the Anterior Pituitary Gland

10

TIME

	BRAIN REGIONS	TIME				
		2 HR	4 HR	8 HR	12 HR	24 HR
15	Medial Basal Hypothalamus	116 \pm 15	154 \pm 24*	110 \pm 20	265 \pm 15*	108 \pm 7
	Preoptic Area	132 \pm 22*	221 \pm 62*	80 \pm 11	67 \pm 4*	98 \pm 15
20	Cerebral Cortex	76 \pm 10*	98 \pm 9	120 \pm 17	215 \pm 28*	100 \pm 7
	Hippocampus	132 \pm 18*	125 \pm 14*	71 \pm 10*	88 \pm 6	93 \pm 5
	Striatum	94 \pm 30	140 \pm 17*	62 \pm 9*	88 \pm 6	103 \pm 7
	Cerebellum	121 \pm 29	120 \pm 14*	73 \pm 9*	110 \pm 62	107 \pm 9
	Brainstem	111 \pm 24	90 \pm 7	70 \pm 6	118 \pm 12	94 \pm 11
25	Anterior Pituitary	95 \pm 10	220 \pm 39*	142 \pm 28	66 \pm 6*	125 \pm 7*

* p < 0.05 versus control value within a brain region. Evaluation of the time effect was made with a one-way ANOVA. After analysis and for clarity of presentation data was transformed to percent of mean control.

30

- 44 -

TABLE XI

**Estrogen and Glucose Uptake in Various Brain Region
Dose-Dependent Effects of 4 Hour E₂B Exposure on Glucose Uptake
in a Variety of Brain Regions and the Anterior Pituitary Gland**

10	BRAIN REGIONS	E ₂ B DOSE		
		1 µg/Kg	10 µg/Kg	100 µg/Kg
15	Medial Basal Hypothalamus	110 ± 23	154 ± 24*	192 ± 28*
	Preoptic Area	98 ± 45	221 ± 62*	155 ± 47*
	Cerebral Cortex	95 ± 27	98 ± 9	87 ± 11
20	Hippocampus	90 ± 19	125 ± 14*	87 ± 12
	Striatum	111 ± 29	140 ± 17*	74 ± 8*
	Cerebellum	112 ± 20	120 ± 14*	57 ± 11*
	Brainstem	116 ± 22	90 ± 7	47 ± 16
25	Anterior Pituitary	112 ± 32	220 ± 39*	64 ± 25

30 * p < 0.05 versus control value. Evaluation of the dose effect was accomplished with a one-way ANOVA. Post hoc comparisons were done with a Dunnett's test. After analysis and for clarity of presentation data was transformed to percent of mean control.

- 45 -

TABLE XII

Estrogen and Glucose Uptake in Various Brain Regions

5

Effect of E₂B on the Extraction of
C¹⁴-2-Deoxyglucose and ³H₂O in the Brain

10

15

GROUP	E ₂ DG	E _{3H2O}
Oil	0.0803 ± 0.011	0.1318 ± 0.943
E ₂ B	0.1208 ± 0.013*	0.1486 ± 0.313


20

* p < 0.05 when compared to oil control. Analyzed with
a t-test for independent samples; n = 7, mean ± sem.

- 46 -

We claim:

1. A method of protecting a population of nerve cells from death, comprising: administering to a nerve cell population in an animal subject, an effective dosage of an estrogen compound sufficient to cause the nerve cell population to be protected from progressive cell damage leading to the death of the cells otherwise occurring without any intervention, the compound having a general structure:



a tautomer thereof, or a pharmaceutically acceptable salt thereof.

- 20 2. A method according to claim 1, further comprising the R2-group on C17 being an hydroxyl in an α isomeric position, and R1 on C3 being a hydroxyl in a β isomeric position.
- 25 3. A method according to claim 1, further comprising the R2-group on the C17 being an hydroxyl group in a β isomeric position, and R1-group on the C3 being an hydroxyl in the β position.
- 30 4. A method according to claim 1, wherein the R1 group and the R2 group are independently selected from the group consisting of hydrogen, hydroxyl, oxygen, methyl, methyl ester, acetate, ethyl ether, 3,3 dimethyl ketal, 17,17 dimethyl ketal, ethynyl- α , benzoate, benzyl ether,
- 35 glucuronide, valerate, cyclopentylpropionate, propionate, hemisuccinate, palmitate, nanthate, stearate, cypionate.

- 47 -

5. A method according to claim 1, wherein the R1 group and the R2 group are independently selected from the group consisting of sodium glucuronide, sulfite sodium salt, sodium phosphate, and trimethyl ammonium salt.

5

6. A method according to claim 1, wherein the cell population includes neuronal cells.

7. A method according to claim 6, wherein the neuronal
10 cells include cholinergic neurons.

8. A method according to claim 6, wherein the neuronal cells include hippocampal cells.

15 9. A method according to claim 6, wherein the neuronal cells include cortex cells.

10. A method according to claim 9, wherein the neuronal cells include glial cells.

20

11. A method according to claim 1, wherein the step of administering an α estrogen compound to a nerve cell population is preceded by the step of administering the compound to a site within a subject so as to decrease loss
25 of neuronal activity in the subject.

12. A method according to claim 11, wherein the site of loss of neuron activity in the subject is selected from the group consisting of hippocampus, cortex and basal
30 forebrain.

13. A method according to claim 11, wherein the loss of neuronal activity is associated with a chronic degenerative disorder.

35

- 48 -

14. A method according to claim 11, wherein the loss of neuronal activity is associated with an acute degenerative disorder.

5 15. A method according to claim 11, wherein the loss of neuronal activity is associated with trauma at the site.

16. A method according to claim 11, wherein the loss of neuronal activity is associated with overstimulation of
10 the excitatory amino acid receptors on neurons.

17. A method according to claim 11, further comprising administering the estrogen compound by one of the group of routes consisting of oral, buccal, intramuscular,
15 transdermal, intravenous and subcutaneous.

18. A method according to claim 17, further comprising administering the estrogen compound in a controlled release vehicle.
20

19. A method according to claim 17, further comprising administering the estrogen compound orally.

20. A method according to claim 17, further comprising
25 administering the estrogen compound subcutaneously.

21. A method according to claim 11, further comprising the step of causing a reversal of loss of memory and loss of learning function.
30

22. A method of protecting a population of cells in a subject from cell death, comprising: administering to the subject, an estrogen compound in a dosage sufficient to cause the cell population to be protected from progressive
35 cell damage leading to the death of the cells otherwise occurring without any intervention.

- 49 -

23. A method according to claim 22, wherein the cause of the progressive cell damage leading to the death of cells is selected from a group consisting of a neurodegenerative disorder, trauma, and aging.

5

24. A method according to any of claim 22 or 23, wherein the progressive cell damage is associated with loss of neuronal activity, and the dosage of the estrogen compound is sufficient to allow the compound to decrease the
10 loss of neuronal activity in the subject.

25. A method according to any of claim 22-24 wherein the cause of progressive cell damage includes an overstimulation of excitatory amino acid receptors.

15

26. A method according to claim 25, further comprising the step of reacting the estrogen compound with an excitatory amino acid receptor so as to block receptor activity.

20

27. A method according to any of claims 22-26, wherein the neurodegenerative disorder is a chronic neurodegenerative disorder.

25 28. A method according to any of claims 22-26, wherein the loss of neuronal activity is associated with an acute neurodegenerative disorder.

29. A method according to any of claims 22-24 and 27-
30 28, wherein the dosage of the estrogen compound administered to the subject is sufficient to cause an increase in the amount of neurotrophic growth factor available to the population of cells.

35 30. A method according to claim 29, wherein the neurotrophic factor is nerve growth factor.

- 50 -

31. A method according to claim 29, wherein the neurotrophic factor is brain derived neurotrophic factor.

32. A method according to any of the preceding claims,
5 wherein the cell population is neuronal and the neuronal cells are protected from axonal retrograde degeneration in the presence of the estrogen compound.

33. A method according to any of the preceding claims,
10 wherein the cell population include neurons and the neurons are protected from apoptosis in the presence of the estrogen compound.

34. A method according to any of the preceding claims,
15 wherein the population of cells is located at a site selected from the hippocampus, the cortex, and the basal forebrain.

35. A method according to claim 34, wherein the
20 population of cells are cholinergic neurons.

36. A method according to any of the preceding claims,
wherein the estrogen compound is administered by one of the group of routes consisting of oral, buccal, intramuscular,
25 transdermal, intravenous and subcutaneous.

37. A method according to any of claims 22-35, further comprising administering the estrogen compound orally.

30 38. A method according to any of claims 22-35, further comprising administering the estrogen compound subcutaneously.

39. A method according to any of claims 22-35, further
35 comprising administering the estrogen compound intramuscularly.

- 51 -

40. A method according to any of the preceding claims, wherein the neurons are located within a first site in a brain of a subject and the step of administering an estrogen compound to the neurons further comprises administering the
5 estrogen compound at a second site distal to the neurons.

41. A method according to any of the preceding claims, wherein the estrogen compound is a steroid.

10 42. A method according to any of the preceding claims, wherein the estrogen compound is 17 β -estradiol, the 17 β -estradiol being administered in an amount sufficient to provide a plasma level in the range 20 pg/ml to 250 pg/ml of 17 β -estradiol.

15

43. A method according to claim 41, wherein the estrogen compound is an isomer of 17 β -estradiol.

44. A method according to any of the preceding claims,
20 further comprising the step of measuring the decrease in nerve cell death by monitoring levels of high affinity choline uptake by the cholinergic neurons.

45. A method according to claim 44, further comprising
25 causing an increase in the levels of choline acetyltransferase.

46. A method of protecting a population of cells in a subject from cell death, comprising the invention as
30 described and illustrated in the specification.

1/10




FIG.1

2/10




FIG.2




FIG.3

4/10




FIG. 4

SUBSTITUTE SHEET (RULE 26)

5/10




FIG. 5

SUBSTITUTE SHEET (RULE 26)

6/10




FIG.6

7/10




FIG.7

SUBSTITUTE SHEET (RULE 26)

8/10




FIG. 8A




FIG. 8B

SUBSTITUTE SHEET (RULE 26)

R ₁ AND/OR R ₂ SUBSTITUTIONS	
NAME	STRUCTURE
HYDROXYL	-OH
METHYL	-CH ₃
METHYL ESTER	-OCH ₃
ACETATE	$\begin{array}{c} \text{O}-\text{C}-\text{CH}_3 \\ \parallel \\ \text{O} \end{array}$
ETHYL ETHER	O-CH ₂ -CH ₃
3, 3, (OR 17, 17) DIMETHYL KETAL	$\begin{array}{c} \diagup \text{OCH}_3 \\ \diagdown \text{OCH}_3 \end{array}$
ETHYNYL- α	$\begin{array}{c} \diagup \text{C} \equiv \text{CH} \\ \diagdown \text{OH} \end{array}$
BENZOATE	$\begin{array}{c} \text{O} \\ \parallel \\ \text{O}-\text{C} \\ \\ \text{C}_6\text{H}_5 \end{array}$
BENZYL ETHER	OCH ₂ -C ₆ H ₅
GLUCURONIDE	C ₆ H ₈ O ₆
SULFATE, SODIUM SALT	OSO ₃ Na
OXIDE	=O
VALERATE	-C ₅ H ₉ O
CYCLOPENTYLPROPIONATE	$-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{CH}_2)_2-\text{C}_5\text{H}_9$
PROPIONATE	$-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{CH}_2)_2$
HEMISUCCINATE	-C ₄ H ₄ O ₃
PALMITATE	-C ₁₆ H ₃₂ O ₂




R ₁ AND/OR R ₂ SUBSTITUTIONS	
NAME	STRUCTURE
SODIUM PHOSPHATE	$-O-PO_3Na_2$
ENANTHATE	$-C_7H_{12}O$
GLUCURONIDE, SODIUM SALT	$-C_6H_8O_6Na$
STEARATE	$-C_{18}H_{34}O$
TRIETHYL AMMONIUM SALT	$-N-(C_2H_5)_3$
CYPIONATE	$O=\overset{\overset{O}{\parallel}}{C}-CH_2CH_2-$ 
17-β- ESTRADIOL	
17α ESTRADIOL	

FIG.9B

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 94/12782

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/565

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	INT.J.DEV.NEUROSCI., vol.5, no.4, 1987 pages 305 - 11 * see abstract *	1-46
X	NEUROSCI.LETT., vol.138, no.1, 1992 pages 157 - 60 see page 158, right column, line 4 - line 16	1-46
X	ADV.BEHAV.BIOL., vol.36, 1989 pages 197 - 212 see page 206, line 21 - page 207, line 44 see page 208, line 7 - page 209, line 12	1-46

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

28 February 1995

Date of mailing of the international search report

23.03.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gerli, P

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 94/12782

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 897 389 (AROONSAKUL) 30 January 1990 cited in the application see claims 1,2 ---	1-46
X	METAB.BRAIN RES., vol.3, no.1, 1988 pages 1 - 18 see page 12 - page 14 ---	1-46
X	BRAIN RES., vol.608, no.1, April 1993 pages 95 - 100 see page 99 ---	1-46
X	J.COMP.NEUROL., vol.197, 1981 pages 197 - 205 see page 203, right column, line 48 - line 52 ---	1-46
X	J.STEROID BIOCHEM.MOL.BIOL., vol.41, no.3-8, 1992 pages 633 - 5 see page 633, left column, line 1 - line 4 see page 635, left column, line 36 - right column, line 11 -----	1-46

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 94/ 12782

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-46 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/12782

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4897389	30-01-90	EP-A- 0324037	19-07-89
		US-A- 4898856	06-02-90
		US-A- 4791099	13-12-88
		US-A- 4898857	06-02-90
		US-A- 4727041	23-02-88
		US-A- 4902680	20-02-90
		US-A- 5017470	21-05-91
